

Decoupling heavy particles simultaneously

R.J. Crewther^a, S.D. Bass^{b*}, F.M. Steffens^{cd†} and A.W. Thomas^{ae‡}

^aDepartment of Physics and Centre for the Subatomic Structure of Matter (CSSM), University of Adelaide, SA 5005, Australia

^bHigh Energy Physics Group, Institute for Experimental Physics and Institute for Theoretical Physics, Universität Innsbruck, Technikerstrasse 25, A 6020 Innsbruck, Austria

^cInstituto de Fisica Teorica, Rua Pamplona 145, 01405-900 São Paulo, SP, Brazil

^dNFC - FCBEE - Universidade Presbiteriana Mackenzie, Rua da Consolação 930, 01302-907, São Paulo, SP, Brazil

^eJefferson National Laboratory, 12000 Jefferson Ave., Newport News, VA 23606, USA

The renormalization group is extended to cases where several heavy particles are decoupled at the same time. This involves large logarithms which are *scale-invariant* and so *cannot* be eliminated by a change of renormalization scheme. A set of scale-invariant running couplings, one for each heavy particle, is constructed without reference to intermediate thresholds. The entire heavy-quark correction to the axial charge of the weak neutral current is derived to next-to-leading order, and checked in leading order by evaluating diagrams explicitly. The mechanism for cancelling contributions from the top and bottom quarks in the equal-mass limit is surprisingly non-trivial.

1. SEVERAL LARGE SCALES

When the renormalization group (RG) is applied to a problem involving two or more large masses or momenta, the standard procedure is to deal with each asymptotic scale separately, starting with the largest. However, there are many cases of practical interest where this sequential approach will not do. For example, the top quark, the $W^\pm$ and Z^0 bosons, and presumably the Higgs boson have masses in the same region, so in situations where more than one of these particles sets the scale, or there are momenta of similar magnitude, it is not reasonable to suppose that one of the corresponding logarithms dominates the others. Various techniques for dealing with

this situation are being investigated [1,2,3,4,5].

Our observation [6,7,8] is that the RG specifies simultaneous dependence on asymptotic scales

$$M_1 \geq M_2 \geq \dots \geq M_n \gg \text{fixed scale } \mu, \quad (1)$$

for limits in which their logarithms grow large together:

$$\ln(M_i/\mu)/\ln(M_j/\mu) = \text{fixed}, \quad M_i, M_j \rightarrow \infty. \quad (2)$$

The key new feature is the appearance of large logarithms

$$\ln(M_i/M_j) = \ln(M_i/\mu) - \ln(M_j/\mu) \quad (3)$$

which are *scale-invariant*, i.e. independent of the renormalization scale μ . This precludes any use of the standard method [9,10] where a running coupling evolves from threshold to threshold and where, at each new threshold M_j , large *scale-dependent* logarithms are suppressed by choosing a new scale $\mu \sim M_j$.

*Research supported by the Austrian Science Fund (FWF) (grant M770). The CSSM is thanked for its hospitality.

†Supported by FAPESP (03/10754-0), CNPq (308932/2003-0) and MackPesquisa.

‡Supported by the Australian Research Council and by DOE contract DE-AC05-84ER40150, under which SURA operates Jefferson National Laboratory.




Figure 1. Neutrino scattering from nucleons at low momentum transfers q .

Instead, for each heavy particle h_i (or large momentum Q), we construct a scale-invariant running coupling

$$\alpha_{h_i} = \alpha_{h_i}(\ln(M_1/\mu), \dots, \ln(M_n/\mu)) \quad (4)$$

without reference to intermediate thresholds. All asymptotic dependence on $M_1, \dots, M_n$ is carried by $\{\alpha_{h_i}; i = 1, \dots, n\}$. Coefficients in the Taylor series in $\{\alpha_{h_i}\}$ are determined by matching with the result for the sequential limit

$$M_j \rightarrow \infty \text{ for fixed } M_{j+1}, \dots, M_n, \quad j = 1, 2, \dots \quad (5)$$

These ideas are tested for the classic example [9,11,12] of heavy quarks h being decoupled from the weak neutral axial-vector current

$$J_{\mu 5}^{\text{Weak}} = \frac{1}{2} \left\{ \sum_{q=u,c,t} - \sum_{q=d,s,b} \right\} \bar{q} \gamma_\mu \gamma_5 q \quad (6)$$

at low momentum transfer q (Fig. 1). In particular, we check that our results for simultaneous t, b decoupling are consistent with the sequential and $m_b \sim m_t$ limits.

Results are presented for three-colour quantum chromodynamics (QCD) with modified minimally subtracted ($\overline{\text{MS}}$) renormalization at fixed scale $\bar{\mu}$.

2. DIAGRAMMATIC ANALYSIS

In perturbation theory, mass logarithms are first produced by the two-loop graphs of Fig. 2.




Figure 2. The lowest-order diagrams (two of them when fermion arrows are restored) responsible for mass logarithms induced by the neutral current (6). Gluon propagators are shown as wavy lines. Straight lines denote propagators for heavy (h) and light (ℓ) quarks.

When both t and b are heavy, the result is [9]

$$\langle \bar{t} \gamma_\mu \gamma_5 t - \bar{b} \gamma_\mu \gamma_5 b \rangle_6 = \left(\frac{\alpha_6}{\pi} \right)^2 \ln \frac{m_t}{m_b} \gamma_\mu \gamma_5 + O(m_{t,b}^{-2}, \alpha_6^3). \quad (7)$$

A subscript F refers to the number of flavours ($F = 6$ in this case); thus $\alpha_6 = g_6^2/4\pi^2$ is the $\overline{\text{MS}}$ renormalized strong coupling for 6-flavour QCD. Eq. (7) follows directly from Adler's two-loop renormalization Z -factor [13] for the gauge-invariant axial-vector current in quantum electrodynamics (QED).

At this two-loop level, the manner in which m_t and m_b are taken to infinity is immaterial: the limits can be taken separately or together, and α_6 , α_5 , and α_4 need not be distinguished.

However, we must be more careful when summing all leading-order (LO) logarithms

$$\sim \alpha_F^{r+s+1} \ln^r(m_t/\bar{\mu}) \ln^s(m_b/\bar{\mu}), \text{ integer } r, s \geq 0.$$

For example, if m_t and m_b tend to ∞ together, α_5 *cannot* enter the analysis, and the coupling α_4 of the residual $F = 4$ theory must be held fixed (*not* α_6).

Two types of diagrams have to be taken into account. First, there are the diagrams in Fig. 3 which produce LO logarithms controlled by the RG; these depend on the strong coupling α_6 of the *original* 6-flavour theory. Then the diagrams




Figure 3. Graphs responsible for leading-order mass logarithms in ghost-free gauges. Each gluon propagator is dressed with one-loop gluon and quark bubbles (all six flavours q) plus counterterms; these insertions sum to a geometric series, as shown. Tadpoles and other one-point gluon insertions are renormalized to zero. Not shown are heavy-quark corrections to the light-quark propagator which produce finite corrections to make the residual singlet axial-vector current scale-invariant, as in Eq. (12).




Figure 4. Leading-order matching condition for the f -flavour coupling $\alpha_f = g_f^2/4\pi$ produced by decoupling $6 - f$ heavy quarks h from the six-flavour theory with coupling $\alpha_6 = g_6^2/4\pi$.

of Fig. 4 are needed to obtain the LO matching condition expressing α_6 in terms of the coupling α_f of the *residual* theory with f light flavours.

2.1. Preliminary Calculation

First we check the LO amplitude $\Gamma_{\mu 5}^{(t)}$ for the top quark t to decouple from $\bar{t}\gamma_\mu\gamma_5 t - \bar{b}\gamma_\mu\gamma_5 b$. In this case, the top mass m_t acts as a regulator for the singlet component

$$\frac{1}{5} \sum_{q=\ell} \bar{q}\gamma_\mu\gamma_5 q, \quad \ell = u, d, s, c, b$$

of the current $\bar{b}\gamma_\mu\gamma_5 b$ in the residual 5-flavour theory, with a renormalizing Z -factor produced by the diagrams of Fig. 3 in LO approximation. All dependence on α_6 has to be eliminated in favour of α_5 , the quantity held fixed as $m_t \rightarrow \infty$. So we apply Fig. 4 with $f = 5$,

$$\alpha_5 \stackrel{\text{LO}}{=} \alpha_6 / \{1 - (\alpha_6/3\pi) \ln(m_t/\bar{\mu})\}, \quad (8)$$

and invert to obtain the matching condition

$$\alpha_6 \stackrel{\text{LO}}{=} \alpha_5 / \{1 + (\alpha_5/3\pi) \ln(m_t/\bar{\mu})\}. \quad (9)$$

After some work on the diagrams of Fig. 3, we obtain an asymptotic formula

$$\Gamma_{\mu 5} \sim -\langle \bar{b}\gamma_\mu\gamma_5 b \rangle_5 - (6/23)^2 \gamma_\mu\gamma_5 / \ln(m_t/\bar{\mu}) \quad (10)$$

in complete agreement with the result of standard RG analysis [9,12]. The number 23 comes from the gluon self-energy insertions, i.e. from the one-loop factor $33 - 2f$ in the beta function

$$\beta_f(x) = -\frac{x^2}{6\pi}(33 - 2f) - \frac{x^3}{12\pi^2}(153 - 19f) + O(x^4) \quad (11)$$

of the *residual* $f = 5$ theory. The bare vertex $\gamma_\mu\gamma_5$ corresponds to the gauge-invariant singlet

operator normalised scale-invariantly [12,6],

$$\left(J_{\mu 5}^{\text{inv}}\right)_f = \sum_{q=\ell} (\bar{q}\gamma_\mu\gamma_5 q)_f^{\text{g. inv}} \exp \int_0^{\alpha_f} dx \frac{\gamma_f(x)}{\beta_f(x)},$$

$$\ell = \text{all } f \text{ light flavours}, \quad (12)$$

where $f = 5$ in Eq. (10), and where [14,12]

$$\gamma_f(x) = \frac{x^2}{\pi^2}f + \frac{x^3}{36\pi^3}(177-2f)f + O(x^4) \quad (13)$$

is the Callan-Symanzik function for the anomalous dimension of the singlet current.

2.2. A Paradox?

Suppose that both t and b are decoupled from $\bar{t}\gamma_\mu\gamma_5 t - \bar{b}\gamma_\mu\gamma_5 b$. The residual theory has $f = 4$ flavours, and all diagrams in Fig. 3 shrink to the bare coupling $\gamma_\mu\gamma_5$, which corresponds to the $f = 4$ version of the invariant current (12).

This case is interesting because of the role played by the equal-mass limit $m_b = m_t$, for which the answer to any order must vanish. To cancel the t term in (10) at $m_b = m_t$, a *five*-flavour b contribution has been suggested [9,12]:

$$\Gamma_{\mu 5}^{\text{cancel}} \sim (6/23)^2 \gamma_\mu\gamma_5 \times \left\{ (\ln(m_b/\bar{\mu}))^{-1} - (\ln(m_t/\bar{\mu}))^{-1} \right\}. \quad (14)$$

However *sequential* decoupling of t and then b does not reproduce this result. When b is decoupled directly from $\langle \bar{b}\gamma_\mu\gamma_5 b \rangle_5$ in Eq. (10), standard RG theory gives [6]

$$\Gamma_{\mu 5}^{\text{seq}} \sim (36/23) \gamma_\mu\gamma_5 \times \left\{ (25 \ln(m_b/\bar{\mu}))^{-1} - (23 \ln(m_t/\bar{\mu}))^{-1} \right\}. \quad (15)$$

A pure five-flavour result is *not* obtained because the residual theory has *four* flavours – hence the factor 25 ($= 33 - 2f$ for $f = 4$) in the b term.

In fact, there is no paradox here: the limits $m_t \sim m_b$ and $m_t \gg m_b$ are extreme cases of the general limit (2). To understand what is happening, it is necessary to keep track of dependence on the scale-invariant logarithm $\ln(m_t/m_b)$.

2.3. Simultaneous t, b Decoupling

The leading logarithms of diagrams in Fig. 3 do not depend on external momenta or light-quark

masses m_ℓ , so we set them equal to zero. The amplitude can be written as a Euclidean integral over the momentum Q carried by each dressed gluon propagator, with self-energy insertions $\Pi(-Q^2)$:

$$\Gamma_{\mu 5}^{t,b} \Big|_{\text{LO}} = \frac{\alpha_6^2}{2\pi^2} \gamma_\mu\gamma_5 \int_0^1 ds \int_{m_b^2}^\infty \frac{dQ^2}{Q^2 \{1 - \Pi(-Q^2)\}^2} \times \left[\{1 + s(1-s)(Q/m_h)^2\}^{-1} \right]_{h=b}^t. \quad (16)$$

The gluon integral has been restricted to the relevant asymptotic region $|Q| \gtrsim m_b$. It can be further restricted by noting that, as t and b decouple, the factor in square brackets tends to 1 for $m_b \lesssim |Q| \lesssim m_t$ and is negligible otherwise:

$$\Gamma_{\mu 5}^{t,b} \Big|_{\text{LO}} = \frac{\alpha_6^2}{2\pi^2} \gamma_\mu\gamma_5 \int_{m_b^2}^{m_t^2} \frac{dQ^2}{Q^2} \frac{1}{(1 - \Pi(-Q^2))^2}. \quad (17)$$

In this Q -range, the gluon self-energy amplitude is

$$\Pi(-Q^2) \Big|_{\text{LO}} = -\frac{23\alpha_6}{12\pi} \ln \frac{Q^2}{\bar{\mu}^2} + \frac{\alpha_6}{6\pi} \ln \frac{m_t^2}{\bar{\mu}^2}, \quad (18)$$

so we are left with a simple logarithmic integral:

$$\Gamma_{\mu 5}^{t,b} \Big|_{\text{LO}} = \frac{6\alpha_6}{23\pi} \gamma_\mu\gamma_5 \left[- \left\{ 1 + \frac{\alpha_6}{6\pi} \left(21 \ln \frac{m_t}{\bar{\mu}} \right) \right\}^{-1} + \left\{ 1 + \frac{\alpha_6}{6\pi} \left(23 \ln \frac{m_b}{\bar{\mu}} - 2 \ln \frac{m_t}{\bar{\mu}} \right) \right\}^{-1} \right]. \quad (19)$$

The last step is to match α_6 directly to α_4 (Fig. 4 for $h = t, b$)

$$\alpha_6 \Big|_{\text{LO}} = \alpha_4 / \left[1 + (\alpha_4/3\pi) \{ \ln(m_b/\bar{\mu}) + \ln(m_t/\bar{\mu}) \} \right], \quad (20)$$

and eliminate α_6 from (19). This yields the complete set of LO logarithms

$$\Gamma_{\mu 5}^{t,b} \Big|_{\text{LO}} = \frac{6\alpha_4}{23\pi} \gamma_\mu\gamma_5 \left[\left\{ 1 + \frac{\alpha_4}{6\pi} \left(25 \ln \frac{m_b}{\bar{\mu}} \right) \right\}^{-1} - \left\{ 1 + \frac{\alpha_4}{6\pi} \left(23 \ln \frac{m_t}{\bar{\mu}} + 2 \ln \frac{m_b}{\bar{\mu}} \right) \right\}^{-1} \right], \quad (21)$$

and hence the general asymptotic result [8]

$$\Gamma_{\mu 5}^{t,b} \sim \frac{36}{23} \gamma_\mu\gamma_5 \left[\left\{ 25 \ln \frac{m_b}{\bar{\mu}} \right\}^{-1} - \left\{ 23 \ln \frac{m_t}{\bar{\mu}} + 2 \ln \frac{m_b}{\bar{\mu}} \right\}^{-1} \right]. \quad (22)$$

This answer replaces (14) and covers all cases:

1. It *vanishes* for $m_t = m_b$.
2. It reproduces the correct sequential result (15) when t is decoupled and then b .
3. It works for intermediate cases as well:

$$1 < \ln(m_t/\bar{\mu})/\ln(m_b/\bar{\mu}) < \infty$$

3. GENERALIZED RG

Beyond LO, it is very difficult to identify all contributions “by hand”, so the RG becomes essential. Our aim [6,7,8] is to extend the RG to take all large scale-invariant logarithms into account, and so generate asymptotic expansions for the simultaneous limit (2): LO, NLO (next to leading order), NNLO (next to NLO), and so on. Since no choice of scale can eliminate all large logarithms, we simply fix the $\overline{\text{MS}}$ scale $\bar{\mu}$ as $\{m_h\} \rightarrow \infty$, as Witten [15] did for a single heavy quark.

Witten has all asymptotic dependence on m_h carried by a RG invariant running coupling $\tilde{\alpha}_h$. For (say) t decoupled from six-flavour QCD, $\tilde{\alpha}_t$ is defined by the equation

$$\ln(m_t/\bar{\mu}) = \int_{\alpha_6}^{\tilde{\alpha}_t} dx \{1 - \delta_6(x)\}/\beta_6(x), \quad (23)$$

where δ_F denotes the F -flavour Callan-Symanzik function for mass renormalization:

$$\delta_F(x) = -2x/\pi + O(x^2). \quad (24)$$

Then α_6 is eliminated by a matching condition [16] relating it to the residual coupling α_5 which is held fixed as $m_t \rightarrow \infty$; (similarly, light-quark masses have to be matched). This sequence can be continued by decoupling the b quark

$$\ln(m_{b5}/\bar{\mu}) = \int_{\alpha_5}^{\tilde{\alpha}_b} dx \{1 - \delta_5(x)\}/\beta_5(x) \quad (25)$$

with α_5 matched to α_4 , and so on.

We begin by reformulating the matching procedure to make RG invariance manifest. Once

again, let t be the heavy quark, and let $\bar{m}_t$ be its scale-invariant mass [15]:

$$\ln(\bar{m}_t/\bar{\mu}) = \int_{\alpha_6}^{\tilde{\alpha}_h} dx/\beta_6(x). \quad (26)$$

Then RG invariance implies the existence of a *matching function* $\mathcal{F}$ defined by [7]

$$\ln(\bar{m}_t/\bar{\mu}) = \int_{\alpha_5}^{\tilde{\alpha}_t} dx/\beta_5(x) + \mathcal{F}_{6 \rightarrow 5}(\tilde{\alpha}_t). \quad (27)$$

The result of eliminating $\tilde{\alpha}_t$ from Eqs. (26) and (27) is the all-orders matching condition relating α_6 to α_5 . Similarly, there is a matching function $\mathcal{G}$ for light-quark masses. For sequential decoupling, there corresponds a sequence of matching functions $(\mathcal{F}_{6 \rightarrow 5}, \mathcal{G}_{6 \rightarrow 5})$, $(\mathcal{F}_{5 \rightarrow 4}, \mathcal{G}_{5 \rightarrow 4})$ and so on. The $\mathcal{F}$ and $\mathcal{G}$ functions are NNLO and NNNLO respectively.

Now consider a simultaneous limit, e.g. the decoupling of t, b . We seek scale-invariant running couplings α_t and α_b which are defined *without* reference to quantities α_5 and m_{b5} associated with a b threshold, but which tend to Witten’s couplings $\tilde{\alpha}_t$ and $\tilde{\alpha}_b$ in the sequential limit (5).

Our prescription [8] is to start with Witten’s formula (23) for the heaviest quark, and then use scale-invariant logarithms to generate the others. For t, b decoupling, we define α_t and α_b as follows,

$$\begin{aligned} \ln(m_t/\bar{\mu}) &= \int_{\alpha_6}^{\alpha_t} dx \{1 - \delta_6(x)\}/\beta_6(x), \\ \ln(m_t/m_b) &= \int_{\alpha_b}^{\alpha_t} dx \{1 - \delta_5(x)\}/\beta_5(x), \end{aligned} \quad (28)$$

where both m_t and m_b are renormalized masses in the original *six*-flavour theory. The integrands of Eq. (28) are chosen such that, in the sequential limit, α_t and α_b agree with $\tilde{\alpha}_t$ and $\tilde{\alpha}_b$ to NLO accuracy.

The definitions of scale-invariant mass and matching functions also have to be generalised. We define invariant masses $\bar{m}_t$ and $\bar{m}_b$ for the simultaneous limit by the formulas

$$\begin{aligned} \ln(\bar{m}_t/\bar{\mu}) &= \int_{\alpha_6}^{\alpha_t} dx/\beta_6(x), \\ \ln(\bar{m}_t/\bar{m}_b) &= \int_{\alpha_b}^{\alpha_t} dx/\beta_5(x), \end{aligned} \quad (29)$$

and then construct matching functions $\mathcal{F}$ and $\mathcal{G}$ for $6 \rightarrow 4$ flavours, e.g.

$$\ln(\overline{m}_b/\bar{\mu}) = \int_{\alpha_4}^{\alpha_b} dx/\beta_4(x) + \mathcal{F}_{6 \rightarrow 4}(\alpha_t, \alpha_b). \quad (30)$$

The matching condition between α_6 and α_4 in the simultaneous t, b limit is found by eliminating α_t and α_b from Eqs. (29) and (30).

All $\{m_h\}$ dependence in the leading asymptotic power of an amplitude is carried by the running couplings $\alpha_t, \alpha_b, \dots$. Thus, a power series expansion in $\alpha_t, \alpha_b, \dots$ yields coefficient amplitudes of the *residual* theory:

$$\text{Amplitude} = \sum_{k\ell\dots} \alpha_t^k \alpha_b^\ell \dots \mathcal{A}_{k\ell\dots}^{\text{res}}$$

This expansion remains valid for the sequential limit with the *same* coefficient amplitudes (to NLO — beyond that, there are calculable corrections due to $\mathcal{F}$ and $\mathcal{G}$ functions). Hence these coefficients can be found by matching against the result for the sequential limit.

For t, b decoupling from the weak neutral current, we found the result [6,8]

$$\begin{aligned} \bar{t}\gamma_\mu\gamma_5 t - \bar{b}\gamma_\mu\gamma_5 b &=_{\text{NLO}} \frac{6}{23\pi}(\alpha_b - \alpha_t) \\ &\times \left\{ 1 + \frac{125663}{82800\pi}\alpha_b + \frac{6167}{3312\pi}\alpha_t \right\} \left(J_{\mu 5}^{\text{inv}} \right)_4. \end{aligned} \quad (31)$$

Notice that Eq. (28) implies $\alpha_t = \alpha_b$ for $m_t = m_b$, so both LO and NLO terms vanish in this case, as they should. Our definitions imply

$$\begin{aligned} \frac{6}{23\pi}(\alpha_b - \alpha_t) &\sim \frac{36}{23} \left[\left\{ 25 \ln \frac{m_b}{\bar{\mu}} \right\}^{-1} \right. \\ &\quad \left. - \left\{ 23 \ln \frac{m_t}{\bar{\mu}} + 2 \ln \frac{m_b}{\bar{\mu}} \right\}^{-1} \right], \end{aligned} \quad (32)$$

so the LO term in (31) correctly reproduces the diagrammatic result (22).

The complete NLO result for the simultaneous decoupling of t, b, c from $J_{\mu 5}^{\text{Weak}}$ is given at the end of [6]. In that case, the definition

$$\ln(m_b/m_c) = \int_{\alpha_c}^{\alpha_b} dx \{1 - \delta_4(x)\}/\beta_4(x) \quad (33)$$

is added to Eq. (28), and Eqs. (29) and (30) are similarly amended so that α_6 is matched directly to α_3 .

REFERENCES

1. F.V. Tkachov, Phys. Lett. B 412 (1997) 350; M. Beneke and V.A. Smirnov, Nucl. Phys. B 522 (1998) 321; V.A. Smirnov, Applied Asymptotic Expansions in Momenta and Masses, Springer, Berlin & Heidelberg, 2002.
2. J. Blumlein and W.L. van Neerven, Phys. Lett. B 450 (1999) 417.
3. W. Beenakker, S. Dittmaier, M. Krämer, B. Plümper, M. Spira, and P.M. Zerwas, Phys. Rev. Lett. 87 (2001) 201805; L. Reina, S. Dawson and D. Wackeroth, Phys. Rev. D 65 (2002) 053017; H. Eberl, M. Kincel, W. Majerotto and Y. Yamada, Nucl. Phys. B 625 (2002) 372.
4. A. Denner, JHEP Proc. PrHEP hep2001/129, hep-ph/0110155; S. Pozzorini, Nucl. Phys. B 692 (2004) 135.
5. S.J. Brodsky, M.S. Gill, M. Melles, and J. Rathsmann, Phys. Rev. D 58 (1998) 116006; M. Binger and S.J. Brodsky, Phys. Rev. D 69 (2004) 095007.
6. S.D. Bass, R.J. Crewther, F.M. Steffens, and A.W. Thomas, Phys. Rev. D 66 (2002) 031901(R).
7. S.D. Bass, R.J. Crewther, F.M. Steffens, and A.W. Thomas, Phys. Rev. D 68 (2003) 096005.
8. S.D. Bass, R.J. Crewther, F.M. Steffens, and A.W. Thomas, hep-ph/0507278 to be published in Phys. Lett. B.
9. J. Collins, F. Wilczek, and A. Zee, Phys. Rev. D 18 (1978) 242.
10. W. Marciano, Phys. Rev. D 29 (1984) 580; J.C. Collins, Renormalization: an introduction to renormalization, the renormalization group, and the operator-product expansion, Cambridge University Press, Cambridge UK, 1984.
11. D.B. Kaplan and A.V. Manohar, Nucl. Phys. B 310 (1988) 527.
12. K.G. Chetyrkin and J.H. Kühn, Z. Phys. C 60 (1993) 497.
13. S.L. Adler, Phys. Rev. 177 (1969) 2426.
14. R. Köberle and N.K. Nielsen, Phys. Rev. D 8 (1973) 660; J. Kodaira, Nucl. Phys. B 165 (1980) 129; S.A. Larin, Phys. Lett. B 303

- (1993) 113; 334 (1994) 192.
15. E. Witten, Nucl. Phys. B 104 (1976) 445.
16. W. Bernreuther and W. Wetzel, Nucl. Phys. B 197 (1982) 228; W. Bernreuther, Ann. Phys. (N.Y.) 151 (1983) 127.